




DEMOGRAPHICS

Total population (000)	9,833 (2008)
Total under-five population (000)	1,635 (2008)
Births (000)	392 (2008)
Birth registration (%)	43 (2005)
Under-five mortality rate (per 1000 live births)	146 (2008)
Infant mortality rate (per 1000 live births)	90 (2008)
Neonatal mortality rate (per 1000 live births)	43 (2008)
Total under-five deaths (000)	54 (2008)
Maternal mortality ratio (per 100,000 live births)	910 (2005)
Lifetime risk of maternal death (1 in N)	19 (2005)
Total maternal deaths	3,500 (2005)

Under-five mortality rate


Deaths per 1000 live births




Source: UNICEF, 2009

Causes of under-five deaths, 2008

Globally more than one third of child deaths are attributable to undernutrition



Source: WHO/CHERG 2010



Source: WHO/CHERG 2010


INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

NUTRITION

Stunting prevalence (moderate and severe, %)	40 (2008)	Complementary feeding rate (6-9 months, %)	32 (2008)
Wasting prevalence (moderate and severe, %)	8 (2008)	Low birthweight incidence (%)	12 (2005)

Underweight prevalence


Percent children < 5 years underweight for age*



*Based on 2006 WHO reference population


Exclusive breastfeeding

Percent infants < 6 months exclusively breastfed



Vitamin A supplementation

Percent children 6-59 months receiving two doses of vitamin A during calendar year




Source: UNICEF

MATERNAL AND NEWBORN HEALTH

Adolescent birth rate (births per 1,000 women)	153 (2003)
Unmet need for family planning (%)	21 (2005)
Antenatal visits for woman (4 or more visits, %)	50 (2007)
Intermittent preventive treatment for malaria (%)	3 (2005)
C-section rate (total, urban, rural, %)	2, 5, 2 (2007)
(Minimum target is 5% and maximum target is 15%)	
Early initiation of breastfeeding (within 1 hr of birth, %)	35 (2008)
Postnatal visit for baby (within 2 days for home births, %)	---


Causes of maternal deaths

Regional estimates for Sub-Saharan Africa, 1997-2007



Source: WHO 2010

Coverage along the continuum of care




Source: DHS, MICS, Other NS

*See Annex for indicator definition

Antenatal care


Percent women aged 15-49 years attended at least once by a skilled health provider during pregnancy



Source: DHS, MICS, Other NS

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Source: DHS, MICS, Other NS

Neonatal tetanus protection

Percent of newborns protected against tetanus




Source: WHO/UNICEF

CHILD HEALTH

Immunization


Percent of children immunised against measles
Percent of children immunised with 3 doses DPT
Percent of children immunised with 3 doses Hib



Source: WHO/UNICEF

Malaria prevention


Percent children < 5 years sleeping under ITNs



Source: DHS

Prevention of mother to child transmission of HIV


Percent HIV+ pregnant women receiving ARVs for PMTCT



Source: UNICEF/UNAIDS/WHO

Diarrhoeal disease treatment


Percent children < 5 years with diarrhoea receiving oral rehydration therapy or increased fluids, with continued feeding



Source: MICS, DHS

Malaria treatment


Percent febrile children < 5 years using antimalarials



Source: DHS

Pneumonia treatment

Percent children < 5 years with suspected pneumonia taken to appropriate health provider
Percent children < 5 years with suspected pneumonia receiving antibiotics




Source: DHS, MICS, DHS

WATER AND SANITATION

Water


Percent population using improved drinking water sources



Source: WHO/UNICEF JMP, 2010

Sanitation

Percent population using improved sanitation facilities




Source: WHO/UNICEF JMP, 2010

EQUITY

Coverage gap by wealth quintile

Average of eight key indicators



Source: DHS 2005

POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	No
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	No
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Resources

Per capita total expenditure on health (US\$)	62 (2007)
General government expenditure on health as % of total government expenditure (%)	5 (2007)
Out-of-pocket expenditure as % of total expenditure on health (%)	89 (2007)
Density of health workers (per 10,000 population)	1.4 (2005)
Official Development Assistance to child health per child (US\$)	4 (2007)
Official Development Assistance to maternal and neonatal health per live birth (US\$)	8 (2007)
National availability of Emergency Obstetric Care services (% of recommended minimum)	25 (2002)